

A NOTE ON q -BERNOULLI NUMBERS AND POLYNOMIALS

TAEKYUN KIM

*Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea*

ABSTRACT. By using q -integration, we will give some integral equation which are related to the Barnes' multiple Bernoulli numbers. The object of this paper is to give explicit q -integral's formulae which are related to Barnes' multiple q -Bernoulli polynomials.

§1. INTRODUCTION

Let $\mathbb{Z}$, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will be denoted by the ring of integers, the ring of p -adic integers, the field of p -adic numbers and the completion of algebraic closure of $\mathbb{Q}_p$ and let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$. When one talks of q -extension, q is variously considered as an indeterminate, a complex number or a p -adic number. If $q \in \mathbb{C}_p$, then we normally assume $|q - 1|_p < p^{-\frac{1}{p-1}}$, so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$.

In this paper, we use the notation:

$$[x] = [x : q] = \frac{1 - q^x}{1 - q}.$$

Hence, $\lim_{q \rightarrow 1} [x] = x$ for any x with $|x|_p \leq 1$ in the present p -adic case.

Key words and phrases. p -adic q -integrals, multiple Barnes' Bernoulli numbers.
2000 Mathematics Subject Classification: 11S80, 11B68, 11M99 .

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

Let d be a fixed integer and let p be a fixed prime number. We set

$$\begin{aligned} X &= \varprojlim_N (\mathbb{Z}/dp^N\mathbb{Z}), \\ X^* &= \bigcup_{\substack{0 < a < dp \\ (a,p)=1}} a + dp\mathbb{Z}_p, \\ a + dp^N\mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^N$, cf. [2, 3, 4, 5, 6, 7, 8].

For any positive integer N , we set

$$\mu_q(a + dp^N\mathbb{Z}_p) = \frac{q^a}{[dp^N]}$$

and this can be extended to a distribution on X . This distribution yields an integral for each non-negative integer m :

$$\int_{\mathbb{Z}_p} [x]^m d\mu_q(x) = \int_X [a]^m d\mu_q(a) = \frac{1}{(1-q)^m} \sum_{i=0}^m \binom{m}{i} (-1)^i \frac{i+1}{[i+1]}.$$

The multiple Barnes' Bernoulli polynomials were defined by

$$(1) \quad \left(\prod_{j=1}^r \frac{w_j}{e^{w_j t} - 1} \right) t^r e^{xt} = \sum_{n=0}^{\infty} B_n^{(r)}(x|w_1, w_2, \dots, w_r) \frac{t^n}{n!},$$

for each $w_j > 0, 0 < t < 1$, cf. [1].

The numbers $B_n^{(r)}(w_1, w_2, \dots, w_r) = B_n^{(r)}(0|w_1, w_2, \dots, w_r)$ are called the multiple Barnes' numbers.

Throughout this paper, we assume that $\alpha_1, \dots, \alpha_k$ are taken in the set of integers and let $w \in \mathbb{Z}_p$. Now, we can consider the multiple q -Bernoulli numbers by using q -integral as follows:

$$\beta_n^{(r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r) = \underbrace{\int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [w + \alpha_1 x_1 + \cdots + \alpha_r x_r]^n d\mu_q(x_1) \cdots d\mu_q(x_r),$$

and

$$\beta_n^{(r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) = \underbrace{\int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^n d\mu_q(x_1) \cdots d\mu_q(x_r).$$

In this paper, we construct the numbers $\beta_n^{(h,r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r)$ for $h \in \mathbb{Z}$ which reduce the multiple Barnes' Bernoulli numbers $B_n^{(r)}(\alpha_1, \alpha_2, \dots, \alpha_r)$ as $q \rightarrow 1$. Also, we give the new explicit formulas which are related to these numbers.

2. AN EXTENSION OF CHANGHEE q -BERNOULLI NUMBERS

For $h \in \mathbb{Z}$, we define the extension of Changhee q -Bernoulli polynomials, numbers as follows:

$$\begin{aligned} & \beta_n^{(h,r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r) \\ (2) \quad &= \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} q^{\sum_{i=1}^r (h-i)x_i} [w + \alpha_1 x_1 + \cdots + \alpha_r x_r]^n d\mu_q(x_1) \cdots d\mu_q(x_r), \end{aligned}$$

and

$$\begin{aligned} & \beta_n^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) \\ (3) \quad &= \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^n q^{\sum_{i=1}^r (h-i)x_i} d\mu_q(x_1) \cdots d\mu_q(x_r). \end{aligned}$$

These can be written as

$$\beta_n^{(h,r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r) = \sum_{j=0}^n \binom{n}{j} q^{wj} \beta_j^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) [w]^{n-j}.$$

By (3), we have

$$\begin{aligned} & \beta_n^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) \\ (4) \quad &= \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^n q^{\sum_{i=1}^r (h-i)x_i} d\mu_q(x_1) \cdots d\mu_q(x_r) \\ &= \frac{1}{(1-q)^n} \sum_{j=0}^n \binom{n}{j} (-1)^j \frac{(j\alpha_1 + h)(j\alpha_2 + h - 1) \cdots (j\alpha_r + h - r + 1)}{[j\alpha_1 + h][j\alpha_2 + h - 1] \cdots [j\alpha_r + h - r + 1]}. \end{aligned}$$

Therefore we obtain the following :

Theorem 1. *For any $n \geq 0$, we have*

$$\beta_n^{(h,r)}(w, q | \alpha_1, \alpha_2, \dots, \alpha_r) = \frac{1}{(1-q)^n} \sum_{j=0}^n \binom{n}{j} (-q^w)^j \prod_{l=1}^r \left(\frac{j\alpha_l + h - l + 1}{[j\alpha_l + h - l + 1]} \right).$$

Remark 1. Note that

$$\begin{aligned} \beta_0^{(2,1)}(q|1) &= \frac{2}{[2]}, \quad \beta_1^{(2,1)}(q|1) = \frac{2q+1}{[2][3]}, \quad \beta_2^{(2,1)}(q|1) = \frac{2q^2}{[3][4]}, \\ \beta_3^{(2,1)}(q|1) &= -\frac{q^2(q-1)(2[3]+q)}{[3][4][5]}, \dots, \\ \beta_0^{(h,1)}(q|1) &= \frac{h}{[h]}, \\ \beta_1^{(h,1)}(q|1) &= -\frac{(1+q+\dots+q^{h-1}) + q(1+q+\dots+q^{h-2}) + \dots + q^{h-1}}{[h][h+1]} \dots \\ \beta_0^{(2,2)}(q|1,1) &= \frac{2!}{[2][1]}, \quad \beta_1^{(2,2)}(q|1,1) = -\frac{2(q+2)}{[2][3]}, \\ \beta_2^{(2,2)}(q|1,1) &= -\frac{2((q-1)^2+5q)}{[3][4]}, \dots, \\ \beta_0^{(r,r)}(q|\underbrace{1, \dots, 1}_{r \text{ times}}) &= -\frac{r!}{[r][r-1] \dots [2][1]}, \quad (\text{ cf. [4, 5, 6, 7, 8, 9, 10, 11] }). \end{aligned}$$

Remark 2. By the definition of $\beta_n^{(h,r)}(q | \alpha_1, \alpha_2, \dots, \alpha_r)$, we note that

$$\beta_n^{(h,r)}(0, q | \alpha_1, \alpha_2, \dots, \alpha_r) = \beta_n^{(h,r)}(q | \alpha_1, \alpha_2, \dots, \alpha_r).$$

By (2),(3), it is easy to see that

$$\begin{aligned}
 & \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^n q^{\sum_{i=1}^r (h-i)x_i} d\mu_q(x_1) \cdots d\mu_q(x_r) \\
 &= (q-1) \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^{n+1} q^{\sum_{i=1}^r (h-\alpha_i-i)x_i} d\mu_q(x_1) \cdots d\mu_q(x_r) \\
 &+ \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^n q^{\sum_{i=1}^r (h-\alpha_i-i)x_i} d\mu_q(x_1) \cdots d\mu_q(x_r).
 \end{aligned}$$

Thus we have

$$\beta_m^{(h,r)}(q| \underbrace{1, 1, \dots, 1}_{r \text{ times}}) = (q-1) \beta_m^{(h-1,r)}(q| \underbrace{1, 1, \dots, 1}_{r \text{ times}}) + \beta_m^{(h-1,r)}(q| \underbrace{1, 1, \dots, 1}_{r \text{ times}}).$$

It is easy to see that

$$\begin{aligned}
 & \underbrace{\int_X \cdots \int_X}_{r \text{ times}} [w + \alpha_1 x_1 + \cdots + \alpha_r x_r]^n q^{\sum_{i=1}^r (h-i)x_i} d\mu_q(x_1) \cdots d\mu_q(x_r) \\
 (5) \quad &= [d]^{n-r} \sum_{i_1, i_2, \dots, i_r=0}^{d-1} q^{(i_1+\dots+i_r)h-i_2-2i_3-\dots-(r-1)i_r} \\
 &\times \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} \left[\frac{w + \alpha_1 i_1 + \cdots + \alpha_r i_r}{d} + \alpha_1 x_1 + \cdots + \alpha_r x_r; q^d \right]^n \\
 &\times q^{x_1(h-1)d+\dots+x_r(h-r)d} d\mu_{q^d}(x_1) \cdots d\mu_{q^d}(x_r).
 \end{aligned}$$

From (3), (5), we have the following:

Theorem 2. *For any positive integer n , we have*

$$\begin{aligned}
 \beta_n^{(h,r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r) &= [d]^{n-r} \sum_{i_1, i_2, \dots, i_r=0}^{d-1} q^{(i_1+\dots+i_r)h-i_2-2i_3-\dots-(r-1)i_r} \\
 &\times \beta_n^{(h,r)}\left(\frac{w + \alpha_1 i_1 + \cdots + \alpha_r i_r}{d}, q^d|\alpha_1, \dots, \alpha_r\right).
 \end{aligned}$$

Moreover,

$$\begin{aligned} \beta_n^{(h,r)}(wd, q|\alpha_1, \alpha_2, \dots, \alpha_r) &= [d]^{n-r} \sum_{i_1, i_2, \dots, i_r=0}^{d-1} q^{(i_1+\dots+i_r)h-i_2-2i_3-\dots-(r-1)i_r} \\ &\quad \times \beta_n^{(h,r)}\left(w + \frac{\alpha_1 i_1 + \dots + \alpha_r i_r}{d}, q^d|\alpha_1, \dots, \alpha_r\right). \end{aligned}$$

Remark 3. Note that

$$\lim_{q \rightarrow 1} \beta_n^{(h,r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r) = B_n^{(r)}(w|\alpha_1, \alpha_2, \dots, \alpha_r), \quad (\text{ see Eq. (1) }).$$

Hence, $\beta_n^{(h,r)}(w, q|\alpha_1, \alpha_2, \dots, \alpha_r)$ can be considered by the q -analogue of Barnes' multiple Bernoulli polynomials. Now, we will give the inverse formula of Eq. (4).

Indeed we see

$$\begin{aligned} &\sum_{i=0}^n \binom{n}{i} (q-1)^i \beta_i^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) \\ &= \sum_{i=0}^n \binom{n}{i} (q-1)^i \underbrace{\int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p}}_{r \text{ times}} [\alpha_1 x_1 + \dots + \alpha_r x_r]^i q^{\sum_{j=1}^r (h-j)x_j} d\mu_q(x_1) \dots d\mu_q(x_r) \\ &= \underbrace{\int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p}}_{r \text{ times}} q^{n(\alpha_1 x_1 + \dots + \alpha_r x_r)} q^{\sum_{j=1}^r (h-j)x_j} d\mu_q(x_1) \dots d\mu_q(x_r) \\ &= \frac{(n\alpha_1 + h)(n\alpha_2 + h - 1) \dots (n\alpha_r + h - r + 1)}{[n\alpha_1 + h][n\alpha_2 + h - 1] \dots [n\alpha_r + h - r + 1]}. \end{aligned}$$

Therefore we obtain the following:

Theorem 3. For $h \in \mathbb{Z}_{\geq 0}$, we have

$$\sum_{i=0}^m \binom{m}{i} (q-1)^i \beta_i^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) = \prod_{j=1}^r \left(\frac{m\alpha_j + h - j + 1}{[m\alpha_j + h - j + 1]} \right).$$

Let χ be a Dirichlet character with conductor $d \in \mathbb{Z}_{\geq 0}$. Then we define the generalized Changhee q -Bernoulli numbers as follows: For $m \geq 0$,

$$(6) \quad \beta_{m,\chi}^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) = \underbrace{\int_X \cdots \int_X}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^m q^{\sum_{j=1}^r (h-j)x_j} \left(\prod_{j=1}^r \chi(x_j) \right) d\mu_q(x_1) \cdots d\mu_q(x_r).$$

By simple calculation, we see that

$$(7) \quad \begin{aligned} & \underbrace{\int_X \cdots \int_X}_{r \text{ times}} [\alpha_1 x_1 + \cdots + \alpha_r x_r]^m q^{\sum_{j=1}^r (h-j)x_j} \left(\prod_{j=1}^r \chi(x_j) \right) d\mu_q(x_1) \cdots d\mu_q(x_r) \\ &= [d]^{m-r} \sum_{i_1, \dots, i_r=0}^{d-1} q^{hi_1 + \cdots + (h-r+1)i_r} \left(\prod_{j=1}^r \chi(x_j) \right) \\ & \times \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{r \text{ times}} \left[\frac{\alpha_1 i_1 + \cdots + \alpha_r i_r}{d} + \alpha_1 x_1 + \cdots + \alpha_r x_r; q^d \right]^m \\ & \times q^{\sum_{j=1}^r x_j (h-j)d} d\mu_{q^d}(x_1) \cdots d\mu_{q^d}(x_r). \end{aligned}$$

By (2),(7), we have the following :

Theorem 4. For $h \in \mathbb{Z}_{\geq 0}$, we have

$$\begin{aligned} \beta_{m,\chi}^{(h,r)}(q|\alpha_1, \alpha_2, \dots, \alpha_r) &= [d]^{m-r} \sum_{i_1, \dots, i_r=0}^{d-1} q^{hi_1 + \cdots + (h-r+1)i_r} \\ & \times \left(\prod_{j=1}^r \chi(x_j) \right) \beta_m^{(h,r)}\left(\frac{\alpha_1 i_1 + \cdots + \alpha_r i_r}{d}, q^d | \alpha_1, \alpha_2, \dots, \alpha_r\right). \end{aligned}$$

Remark. By using our formulae in the case of $q = 1$, we can obtain many new formulas which are related to the multiple Barnes' Bernoulli numbers, cf.[1, 2, 4, 8, 10, 11].

ACKNOWLEDGEMENTS: This paper was supported by Korea Research Foundation Grant((KRF-2003-05-C00009) .

REFERENCES

1. E. W. Barnes, *On the theory of the multiple gamma functions*, Trans. Camb. Philos. Soc. **19** (1904), 374-425.
2. M. Cenkci, M. Can, V. Kurt, *p-adic interpolation functions and Kummer type congruences for q-twisted and q-generalized twisted Euler numbers*, Advan. Stud. Contemp. Math. **9(2)** (2004), 203-216.
3. A. Khrennikov, S. Ludkovsky, *Stochastic processes on non-Archimedean spaces with values in non-Archimedean fields*, Advan. Stud. Contemp. Math. **5(1)** (2002), 57-91.
4. T. Kim, *Sums powers of consecutive q-integers*, Advan. Stud. Contemp. Math. **9** (2004), 15-18.
5. T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
6. T. Kim, *An invariant p-adic integral associated with Daehee numbers*, Integral Trans. Special Funct. **13** (2002), 65-69.
7. T. Kim, *On the Sehee integral representation associated with q-Riemann zeta function*, Proc. Jangjeon Math. Soc. **7 no. 2** (2004), 00-00.
8. T. Kim, S. H. Rim, *A note on two-variable Dirichlet's L-functions*, Advan. Stud. Contemp. Math. **10 no. 1** (2004), 1-6.
9. T. Kim, *p-adic q-integrals associated with Changhee-Barnes' q-Bernoulli polynomials*, Integral Trans. Special Funct. **15** (2004), 415-420.
10. C.S. Ryoo, H. Song, R.P. Agarwal, *On the roots of the q-analogue of Euler-Barnes' polynomials*, Advan. Stud. Contemp. Math. **9(2)** (2004), 153-163.
11. M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts, and qunits*, The Electronic J. Combinatorics **11** (2004), R 71.